

Contact information for Hantavirus Training  
Hantavirus orientation by IHS  
Conducted by Sam Canyon, IHS Environmental Health

General information

1. Study the lease agreement; understand the responsibilities of each party (for storage units).
2. Surveillance of the building, check for any holes in the building, check the ceiling, put gravel down around the skirt so rodents cannot dig under the skirt.
3. Food and coffee do not leave out in the open. Pour coffee out at end of the day. Do not leave any food out or in your desk, mice will chew through unopened packages of food.
4. Plants, water in the morning, mice will drink from excess water which settles in the plant container.
5. Mice move to the food source, do not leave food unattended on the desk.
6. Mice will chew through wood, plastic, i.e. telephone pad.
7. Trap mice out to 1000 feet from the building.
8. One hundred percent safety for all employees.
9. Category 4 disease, very dangerous.
10. Hantavirus came from North Korea, Hantan river region.
11. Man does not compete with mice anymore for food, mice population has risen.
12. If you own a cat which hunts and kills mice, do not let it into the house because the fleas from the mouse jump onto the cat.
13. Deer mice droppings, saliva, and urine; attach to particles of dust and become airborne when disturbed which you inhale and become infected.
14. Lungs become full of fluid and you drown in your own fluids.
15. Mouse head is made of cartilage which enables it to squeeze through tiny openings.
16. To kill the hantavirus, spray with a mixture of 90% water and 10% bleach.
17. Spray mouse dropping thoroughly so dropping won't become airborne.
18. Spray Lysol into the air to kill the virus and as an extra precaution.
19. Sunlight will kill hantavirus.
20. Hantavirus is smaller than bacteria, need a very efficient mask to block the tiny hantavirus.
21. Need special mask.
22. A dead mouse will give off vapors, so be very careful when breathing around dead mice.

To Proceed with Cleaning


23. Identify hot zone
24. Identify wind direction, wind at your back.
25. Command post
26. Danger zone

27. Decontamination zone
28. Type 49 HEPA filter
29. Full protective clothing
30. Cover you head and your feet.
31. Throw in a fogger first, bug bomb.
32. Air out storage area for a couple hours first, very important
33. Put equipment in the sunlight which kills the virus.
34. Check storage for holes and plug the holes.
35. Flea spray
36. Safe zone – suit up or take off protective gear.
37. Air out storage area for a few hours and spray
38. Foggers to kill fleas and spiders.
39. Spray the area hard and long
40. Spray Lysol into the air
41. Throw away your protective clothing
42. Contact with chlorine may cause sores or allergies.
43. Food to use with traps, 4 way grain with peanut butter
44. Do not use poison, mouse may die in the wall of your building
45. Do not use glue traps
46. Visit Native American Research website.
47. If you suspect you are infected, get to the hospital immediately.
48. One treatment, pump fresh blood into you.
49. Oxygen to help you breathe.

#### Snakes

50. Rattlesnakes have a five foot strike zone
51. Peppers may be use to kill rattlesnakes, instructor did not specify, you may need to experiment.
52. Kill the rattlesnake old-fashioned way with shovel or any tool you are comfortable using.

Command  
Post  
Zones  
Wind



Hot

Danger

Decon

Safe